


AD 698 800

THE MAXIMUM AND MINIMUM OF A POSITIVE DEFINITE  
QUADRATIC POLYNOMIAL ON A SPHERE ARE CONVEX  
FUNCTIONS OF THE RADIUS

George E. Forsythe

Stanford University  
Stanford, California

July 1969



This document has been approved for public release and sale.

00888631

CS 144

THE MAXIMUM AND MINIMUM OF A POSITIVE DEFINITE  
QUADRATIC POLYNOMIAL ON A SPHERE ARE  
CONVEX FUNCTIONS OF THE RADIUS

BY

GEORGE E. FORSYTHE

TECHNICAL REPORT NO. CS 144  
JULY 1969

CLEARINGHOUSE

COMPUTER SCIENCE DEPARTMENT  
School of Humanities and Sciences  
STANFORD UNIVERSITY



THE MAXIMUM AND MINIMUM OF A POSITIVE DEFINITE  
QUADRATIC POLYNOMIAL ON A SPHERE ARE  
CONVEX FUNCTIONS OF THE RADIUS

by

George E. Forsythe

July 1969

Reproduction in whole or in part is permitted  
for any purpose of the United States Government.

Computer Science Department  
Stanford University

### Abstract

It is proved that in euclidean  $n$ -space the maximum  $M(\rho)$  and minimum  $m(\rho)$  of a fixed positive definite quadratic polynomial  $Q$  on spheres with fixed center are both convex functions of the radius  $\rho$  of the sphere. In the proof, which uses elementary calculus and a result of Forsythe and Golub,  $m''(\rho)$  and  $M'(\rho)$  are shown to exist and lie in the interval  $[2\lambda_1, 2\lambda_n]$ , where  $\lambda_i$  are the eigenvalues of the quadratic form of  $Q$ . Hence  $m''(\rho) > 0$  and  $M'(\rho) > 0$ .

### Summary

Let  $A$  be a given symmetric, nonsingular matrix of real elements and order  $n$ . Let  $b$  be a given column vector of  $n$  real elements. For each real column  $n$ -vector  $x$ , the nonhomogeneous quadratic polynomial

$$Q(x) = (x-b)^T A(x-b)$$

( $T$  denotes transpose) is a real number. Let  $\lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_n$  be the (necessarily) real eigenvalues of  $A$ . Let  $m(\rho)$  be the minimum of  $Q(x)$  on the sphere  $S_\rho = \{x: x^T x = \rho^2\}$ , and let  $M(\rho)$  be the maximum of  $Q(x)$  on  $S_\rho$ . M. J. D. Powell asked the author whether  $m(\rho)$  is a convex function of  $\rho$  when  $A$  is positive definite. An affirmative answer is given by the theorem:

- (1) Theorem. If  $A$  is positive definite (i.e., if  $0 < \lambda_1$ ), then both  $m(\rho)$  and  $M(\rho)$  are convex functions of  $\rho$ , for all  $\rho > 0$ .

Theorem (1) will follow from the following result:

- (2) Theorem. Let  $A$  be any nonsingular matrix. Then for  $\rho > 0$ , the second derivatives  $m''(\rho)$  and  $M''(\rho)$  both exist, and

$$(3) \quad m''(\rho) \geq 2\lambda_1 \quad \text{and} \quad M''(\rho) \geq 2\lambda_1.$$

Equality occurs in (3) if and only if  $Ab = \lambda_1 b$ . Moreover,

$$(4) \quad m''(\rho) \leq 2\lambda_n \quad \text{and} \quad M''(\rho) \leq 2\lambda_n$$

and equality occurs in (4) if and only if  $Ab = \lambda_n b$ .

### Review of Previous Work

The proof of Theorem (2) is based on techniques developed in Forsythe and Golub [1], which dealt only with the case  $\rho = 1$ . The relevant results of [1] are now summarized and extended to general  $\rho$ .

Let  $\{u_1, \dots, u_n\}$  be an orthonormal real set of eigenvectors of  $A$ , with  $Au_i = \lambda_i u_i$  ( $i = 1, \dots, n$ ). Let  $b = \sum b_i u_i$ . For any vector  $x$  in  $S_\rho$  at which  $Q(x)$  is stationary with respect to  $S_\rho$ , there is a real number  $\lambda$  with

$$(5) \quad A(x-b) = \lambda x$$

$$(6) \quad x^T x = \rho^2.$$

Letting  $x = \sum x_i u_i$ , we find from (5) that

$$(7) \quad x_i = \frac{x_i b_i}{\lambda_i - \lambda},$$

so that (6) becomes

$$(8) \quad r(\lambda) \equiv \sum_{i=1}^n \frac{\lambda_i^2 b_i^2}{(\lambda_i - \lambda)^2} = \rho^2.$$

For each given value of  $\rho > 0$ , equation (8) determines from 2 to  $2n$  real values of  $\lambda$ . For each  $\lambda$  so determined, equation (5) determines one or more vectors  $x^\lambda$  (if all  $b_i \neq 0$ , then  $x^\lambda$  is unique). For any  $x^\lambda$ , we have

$$(9) \quad Q(x^\lambda) = r(\lambda),$$

where

$$(10) \quad r(\lambda) = \lambda^2 \sum_{i=1}^n \frac{\lambda_i b_i^2}{(\lambda_i - \lambda)^2} .$$

Now  $Q(x)$  is stationary with respect to  $S_\rho$  at any  $x^\lambda$ . For given  $\rho$ , let  $\Lambda_L = \Lambda_L(\rho)$  and  $\Lambda_R = \Lambda_R(\rho)$  be the smallest resp. largest values of  $\lambda$  satisfying equation (8). Theorem (4.1) of [1] states that  $f(\Lambda_L)$  and  $f(\Lambda_R)$  are the minimum resp. maximum values of  $Q(x)$  on  $S_\rho$ .

Much of [1] was devoted to the singular cases where some  $b_i = 0$ . For the present investigation, where we are interested only in the values of  $Q(x)$ , we simply omit from the sums (8) and (10) all terms with  $b_i = 0$ , and reduce  $n$ , if necessary. Having done that, it is then clear from (8) that, for any  $\rho$ ,

$$(11) \quad \Lambda_L < \lambda_1 \quad \text{and} \quad \lambda_n < \Lambda_R .$$

This concludes the necessary summary of [1].

As a digression, the author notes that the main theorems (2.7) and (4.1) of [1] were proved in [1] by studying  $f(\lambda)$  and  $g(\lambda)$  for complex values of  $\lambda$ . In late 1965, Professor W. Kahan [unpublished] showed us how to prove those theorems more simply, using only real values of  $\lambda$ .

#### Proof of Theorem (2).

With the above apparatus our problem is reduced to an exercise in the differential calculus. For each  $\rho > 0$  we determine a unique Lagrange multiplier  $\lambda = \lambda(\rho)$  from (8) -- either the minimal  $\Lambda_L$  or maximal  $\Lambda_R$ . For ease of exposition, suppose  $\lambda(\rho) = \Lambda_L$ . Then the function

$$(12) \quad m(\rho) = f(\lambda(\rho))$$

is determined from (10). Since  $f(\lambda)$  and  $g(\lambda)$  are analytic for  $\lambda < \lambda_1$ , the function  $m(\rho)$  has derivatives of all order. We shall determine  $m''(\rho)$  by calculus. To simplify some expressions, we introduce the abbreviations

$$(13) \quad \alpha_p = \sum_{i=1}^n \frac{\lambda_i^2 b_i^2}{(\lambda_i - \lambda)^2} \quad (p = 2, 3, 4) .$$

Differentiating (10) and simplifying, we find:

$$(14) \quad \frac{df}{d\lambda} = 2\lambda\alpha_3 ;$$

$$(15) \quad \frac{d^2f}{d\lambda^2} = 2\alpha_3 + 6\lambda\alpha_4 .$$

Now equation (8) states that, when  $\lambda = \lambda(\rho)$ ,

$$(16) \quad \alpha_2 = \rho^2 .$$

Differentiating (8) twice with respect to  $\rho$  yields

$$(17) \quad \frac{d\lambda}{d\rho} \alpha_3 = \rho ;$$

$$(18) \quad \frac{d^2\lambda}{d\rho^2} \alpha_3 + 3\left(\frac{d\lambda}{d\rho}\right)^2 \alpha_4 = 1 .$$

Solving (17) and (18) in turn, we find

$$(19) \quad \frac{d\lambda}{d\rho} = \frac{\rho}{\alpha_3} ;$$

$$(20) \quad \frac{d^2\lambda}{d\rho^2} = \frac{1}{\alpha_3} - \frac{3\rho^2\alpha_4}{\alpha_3^2} .$$



Now, by the chain rule,

$$\frac{dm}{d\rho} = \frac{df}{d\lambda} \cdot \frac{d\lambda}{d\rho} ,$$

and

$$(21) \quad \frac{d^2 m}{d\rho^2} = \frac{d^2 f}{d\lambda^2} \left( \frac{d\lambda}{d\rho} \right)^2 + \frac{df}{d\lambda} \cdot \frac{d^2 \lambda}{d\rho^2} .$$

We now substitute into (21) the expressions (14), (15), (19), and (20).

We find that

$$(22) \quad m''(\rho) = \frac{d^2 m}{d\rho^2} = (2\alpha_3 + 6\lambda\alpha_4) \frac{\rho^2}{\alpha_3^2} + 2\lambda\alpha_3 \left( \frac{1}{\alpha_3} - \frac{3\rho^2\alpha_4}{\alpha_3^3} \right) .$$

Hence

$$\frac{1}{2} m''(\rho) = \lambda + \frac{\rho^2}{\alpha_3} = \frac{1}{\alpha_3} (\lambda\alpha_3 + \alpha_2) , \quad \text{by (16).}$$

Simplifying,

$$\frac{1}{2} m''(\rho) = \frac{1}{\alpha_3} \sum_{i=1}^n \frac{\lambda_i^3 b_i^2}{(\lambda_i - \lambda)^3} , \quad \text{or}$$

$$(23) \quad \frac{1}{2} m''(\rho) = \sum_{i=1}^n \frac{\lambda_i^3 b_i^2}{(\lambda_i - \lambda)^3} \bigg/ \sum_{i=1}^n \frac{\lambda_i^2 b_i^2}{(\lambda_i - \lambda)^3} .$$

Formula (23) is the end of our calculus exercise. In it,  $\lambda$  is determined from solving (8). Note by (11) that the factors  $(\lambda_i - \lambda)^3$  all have the same sign for  $i = 1, 2, \dots, n$ , whether  $\lambda = \lambda_L$  or  $\lambda = \lambda_R$ . Hence  $\frac{1}{2} m''(\rho)$  is a weighted average with positive weights of the  $\{\lambda_i\}$ .

It follows that  $\frac{1}{2} m''(\rho) \geq \lambda_1$ , with equality only when all  $\lambda_i$  in (23) are equal to  $\lambda_1$ , i.e., if  $b_i = 0$  for  $\lambda_i > \lambda_1$ . This proves (3), and (4) is proved analogously. This concludes the proof of Theorem (2).

It would be desirable to have a simple geometrical proof.

#### What if A is singular?

If A is singular, that is, if some  $\lambda_i = 0$ , the situation is somewhat more complicated, just as the case where some  $\lambda_i b_i = 0$  is complicated in [1]. Theorem (2) fails to hold for semidefinite matrices, because  $m''(\rho)$  may not exist for some  $\rho$ , as the following example shows:

(24) Example. For  $n = 2$  let  $Q(x) = (x_2 - 1)^2$ , where  $x = (x_1, x_2)^T$ .

Then

$$m(\rho) = \begin{cases} 1-\rho & , \quad 0 \leq \rho \leq 1 \\ 0 & , \quad 1 \leq \rho < \infty \end{cases}$$

so  $m''(1)$  does not exist.

If  $\lambda_1 = 0$ , the Lagrange multiplier remains at  $\lambda = 0$  for all sufficiently large  $\rho$ .

Theorem (1) can easily be extended to semidefinite matrices by continuity. We have

(25) Theorem. If A is positive semidefinite (i.e., if  $0 \leq \lambda_1$ ),  
then both  $m(\rho)$  and  $M(\rho)$  are convex functions of  $\rho$  for  $\rho > 0$ .

In proof, we note that  $m(\rho)$  and  $M(\rho)$  are continuous functions of the elements of A. If A is semidefinite, it can be approximated by a definite matrix  $A_\epsilon$ , for which  $m_\epsilon$  and  $M_\epsilon$  are convex, with  $\|A - A_\epsilon\| < \epsilon$ . Letting  $\epsilon \rightarrow 0$ , we find that  $m = \lim m_\epsilon$  and  $M = \lim M_\epsilon$  are convex.

#### Reference

- [1] George E. Forsythe and Gene H. Golub, "On the stationary values of a second-degree polynomial on the unit sphere", J. Soc. Indust. Appl. Math., vol. 13 (1965), pp. 1050-1068.

Unclassified

Security Classification

DOCUMENT CONTROL DATA - R & D		
(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)		
1. ORIGINATING ACTIVITY (Corporate author) Computer Science Department Stanford University Stanford, California 94305		2a. REPORT SECURITY CLASSIFICATION Unclassified
		2b. GROUP --
3. REPORT TITLE  THE MAXIMUM AND MINIMUM OF A POSITIVE DEFINITE QUADRATIC POLYNOMIAL ON A SPHERE ARE CONVEX FUNCTIONS OF THE RADIUS		
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) Manuscript for Publication (Technical Report)		
5. AUTHOR(S) (First name, middle initial, last name)  George E. Forsythe		
6. REPORT DATE July 1969	7a. TOTAL NO. OF PAGES 9	7b. NO. OF REFS 1
8a. CONTRACT OR GRANT NO. N00014-67-A-0112-0029	9a. ORIGINATOR'S REPORT NUMBER(S) CS 144	
b. PROJECT NO. NR 044-211	9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report) none	
c.		
d.		
10. DISTRIBUTION STATEMENT  Releasable without limitations on dissemination.		
11. SUPPLEMENTARY NOTES  ---	12. SPONSORING MILITARY ACTIVITY  Office of Naval Research	
13. ABSTRACT  It is proved that in euclidean n-space the maximum $M(\rho)$ and minimum $m(\rho)$ of a fixed positive definite quadratic polynomial $Q$ on spheres with fixed center are both convex functions of the radius $\rho$ of the sphere. In the proof, which uses elementary calculus and a result of Forsythe and Golub, $m''(\rho)$ and $M''(\rho)$ are shown to exist and lie in the interval $[2\lambda_1, 2\lambda_n]$ , where $\lambda_i$ are the eigenvalues of the quadratic form of $Q$ . Hence $m''(\rho) > 0$ and $M''(\rho) > 0$ .		

DD FORM 1 NOV 65 1473

(PAGE 1)

S/N 0101-807-6801

Unclassified

Security Classification

Unclassified

**Security Classification**

14	KEY WORDS	LINK A		LINK B		LINK C	
		ROLE	WT	ROLE	WT	ROLE	WT
	Maximum on a sphere						
	Minimum on a sphere						
	Quadratic function						
	Convex						

Unclassified

---

Security Classification